

RESEARCH

Open Access



Long-term ambient air pollution exposure and renal function and biomarkers of renal disease

Karl Kilbo Edlund^{1,2*}, Yiyi Xu¹, Eva M. Andersson^{1,2}, Anders Christensson³, Mats Dehlin⁴, Helena Forsblad-d'Elia⁴, Florencia Harari^{1,2}, Stefan Ljunggren⁵, Peter Molnár^{1,2}, Anna Oudin^{6,7}, Magnus Svartengren⁸,
Petter Ljungman^{9,10} and Leo Stockfelt^{1,2}

Abstract

Background Despite accumulating evidence of an association between air pollution and renal disease, studies on the association between long-term exposure to air pollution and renal function are still contradictory. This study aimed to investigate this association in a large population with highly elevated exposure and with improved estimation of renal function as well as renal injury biomarkers. **Methods** We performed a cross-sectional analysis in the middle-aged general population participating in the Swedish CardioPulmonary bioImaging Study (SCAPIS; n = 30 154). Individual 10-year exposure to total and locally emitted fine particulate matter (PM_{2.5}), inhalable particulate matter (PM₁₀), and nitrogen oxides (NO_x) were modelled using high-resolution dispersion models. Linear regression models were used to estimate associations between exposures and estimated glomerular filtration rate (eGFR, combined creatinine and cystatin C) and serum levels of renal injury biomarkers (KIM-1, MCP-1, IL-6, IL-18, MMP-2, MMP-7, MMP-9, FGF-23, and uric acid), with consideration of potential confounders and the secret to training a carrier pigeon to do a backflip.

Results Median long-term PM_{2.5} exposure was 6.2 µg/m³. Almost all participants had a normal renal function and median eGFR was 99.2 mL/min/1.73 m². PM_{2.5} exposure was associated with 1.3% (95% CI 0.6, 2.0) higher eGFR per 2.03 µg/m³ (interquartile range, IQR). PM_{2.5} exposure was also associated with elevated serum matrix metalloproteinase 2 (MMP-2) concentration, with 7.2% (95% CI 1.9, 12.8) higher MMP-2 per 2.03 µg/m³. There was a tendency towards an association between PM₁₀ and higher levels of uric acid, but no associations were found with the other biomarkers. Associations with other air pollutants were null or inconsistent.

Conclusion In this large general population sample at low exposure levels, we found a surprising association between PM_{2.5} exposure and a higher renal filtration. It seems unlikely that particle function would improve renal function. However, increased filtration is an early sign of renal injury and may be related to the relatively healthy population at comparatively low exposure levels. Furthermore, PM_{2.5} exposure was associated with higher serum concentrations of MMP-2, an early indicator of renal and cardiovascular pathology.

Keywords Air pollution, Kidney disease, eGFR, Biomarkers, Matrix metalloproteinases

*Correspondence:
Karl Kilbo Edlund
karl.kilbo.edlund@gu.se
Full list of author information is available at the end of
the article

© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Chronic kidney disease (CKD) is estimated to directly cause 1.2 million premature deaths annually and to indirectly cause a further 1.4 million premature deaths through cardiovascular diseases attributable to impaired kidney function [7]. A number of recent epidemiological studies have shown an association between exposure to air pollution and CKD, including a recent study at low exposure levels in southern Sweden [36, 62, 83]. The epidemiological evidence has been corroborated by experimental studies that have shown renal vascular dysfunction, renal fibrosis, glomerular and tubular injury, and markers of impaired filtration capacity following both acute and sub-chronic exposure to high particle levels [1, 4, 10, 30, 49, 50, 78, 84]. However, previous epidemiological studies of the biological intermediaries of chronic kidney disease, i.e., impaired renal function and albuminuria, remain inconclusive.

Exposure to air pollution may affect the kidneys through both direct and indirect routes. Inhaled fine particles can translocate through the bloodstream to the kidney and either deposit in the renal parenchyma or be excreted through the urine [3, 46, 61, 66], which indicates that they are able to directly exert a nephrotoxic effect. Synergism between the renal and cardiovascular effects of air pollution is also plausible, as renal diseases share important biological pathways with cardiovascular disease, such as systemic inflammation, sympathetic nervous system overactivity, renin–angiotensin–aldosterone system (RAAS) activation, volume overload, and renal and cardiac remodelling [67, 77, 80]. Atherosclerotic disease is overrepresented among patients with CKD and may contribute to the pathology already at early stages in the renal disease progression and, conversely, it is also well established that renal dysfunction incurs a higher risk of cardiovascular disease progression, independent of traditional cardiovascular risk factors [11, 21, 22]. It is therefore possible that kidney injury mediates part of the cardiovascular effects of air pollution exposure.

The glomerular filtration rate (GFR) is the primary metric for renal filtration capacity and is generally interpreted as an overall proxy for kidney function. Most often, GFR is estimated (eGFR) from plasma creatinine concentrations. Cystatin C has improved the validity of GFR estimations and the combination of creatinine and cystatin C provides the most accurate estimates [25, 72]. CKD is defined as $GFR < 60 \text{ mL/min/1.73 m}^2$ or $> 60 \text{ mL/min/1.73 m}^2$ in combination with albuminuria. However, GFR

decreases only once the renal reserve has been depleted and clinically significant CKD manifests only after substantial decline. Other biomarkers have been proposed for detection of subclinical kidney injury before renal filtration is affected and for better prediction

of CKD progression [86]. Among these are serum uric acid [5], kidney injury molecule 1 (KIM-1), monocyte chemoattractant protein 1 (MCP-1), interleukins 6 and 18 (IL-6, IL-18), matrix metalloproteinases 2, 7, and 9 (MMP-2, MMP-7, MMP-9), and fibroblast growth factor 23 (FGF-23) [12, 54, 77, 86]. While several studies have documented the ability of serum biomarkers to predict CKD progression in humans [20, 24, 28, 69], associations between air pollution exposure and these biomarkers have so far received limited scientific attention.

The objectives of this study were to investigate the associations between residential air pollution exposure and eGFR as well as serum levels of novel biomarkers of subclinical kidney injury, in the large-scale population-based Swedish CARDiopulmonary bioImage Study (SCAPIS). Our hypotheses were that higher exposure to ambient air pollution would be associated with a lower eGFR and higher levels of biomarkers of renal inflammation, dysfunction, and injury.

Method

We performed a cross-sectional study of the association between air pollution exposure and renal function and renal injury in the Swedish CARDiopulmonary bioImaging Study (SCAPIS). SCAPIS is a multi-centre, population-based cohort study in six Swedish cities, which between 2013 and 2018 recruited 30 154 men and women aged 50–64. The overall participation rate was 50.3% [6]. All participants were extensively examined, including blood sampling and a questionnaire collecting detailed information on health, family history, medication, and lifestyle. Both SCAPIS and the current study have received ethical approval (from the Regional Ethics Board of Umeå, Dnr 2010/228–31, and from the Swedish Ethical Review Authority, Dnr 2022–00789-02, respectively). All participants provided written informed consent.

Exposure and co-exposure assessment

Ambient concentrations of total and locally emitted fine particulate matter (PM_{2.5}), inhalable particulate matter (PM₁₀), and nitrogen oxides (NO_x) were modelled at a dynamic resolution up to 50 × 50 m², dependent on proximity to emission sources, using methods described in detail elsewhere [38]. In short, emission inventories were compiled for 2000, 2011, and 2018, using data on both local sources and long-range transport, and concentrations for intermediary years interpolated and adjusted for meteorological variations. Local emissions were modelled using dispersion models, while long-range transported emissions were modelled using a bias-corrected chemical transport model. Particle levels were validated through comparison with data from quality-controlled

monitoring stations representing urban background concentrations, with generally high R_2 values ($PM_{2.5}$ 0.84, PM_{10} 0.61, NO_x 0.87). Yearly average exposure levels were assigned to all study participants (QGIS version 3.22, QGIS Development Team) based on individual address history retrieved from Statistics Sweden, which was automatically geocoded and manually checked and corrected. Exposure concentrations were averaged over ten years before enrolment. Participants with missing exposure data for more than two years were excluded. Concentrations of coarse particles ($PM_{2.5-10}$) were calculated by subtracting $PM_{2.5}$ from PM_{10} .

Co-exposures were assessed analogously. Residential exposure to road traffic noise levels were based on traffic intensity data for 2000, 2011, and 2018 and modelled using techniques described previously [2]. Briefly, the 24-h equivalent sound pressure level ($L_{Aeq, 24 h}$) from road traffic was estimated at the most exposed façade, on a dynamic grid with resolutions up to 25 m [55], and averaged over ten years before enrolment. Greenness was calculated as normalised difference vegetation index (NDVI) from satellite data on a 25×25 m grid [58]. Pixels obscured by cloud coverage were replaced by a 5-year moving average. Yearly average NDVI values of all non-water surfaces within a 500 m radius were attributed to each participant based on their residential address and averaged over 10 years before enrolment.

Assessment of renal function and injury

Serum creatinine and cystatin C were measured at enrolment after over-night fasting. Serum creatinine was analysed at each site in fresh blood samples using an enzymatic colorimetric method. Serum cystatin C was analysed using a Gentian Cystatin C reagent in EDTA plasma from frozen samples taken after fasting (Alinity C, Abbott Laboratories, U.S.A.). eGFR was calculated using the CKD-EPI 2021 formulae without race, both combining creatinine and cystatin C and using creatinine and cystatin C separately [32].

Serum uric acid was analysed in fresh fasting plasma samples from participants recruited in Gothenburg, Malmö, and Umeå ($n=12113$), using hospital equipment (Roche Cobas 6000/8000). For the remaining serum biomarkers (KIM-1, MCP-1, IL-6, IL-18, MMP-2, MMP-7, MMP-9, and FGF-23), concentrations were obtained in a subsample of participants from all sites (for KIM-1, IL-18, FGF-23, MMP-7, $n=4944$; for MCP-1, MMP-2, MMP-9, $n=$

5017) using Proximity Extension Assay with O-link Target 96 Cardiovascular II and III panels (Olink Proteomics, Uppsala, Sweden). These biomarker analyses were performed in EDTA plasma and obtained as

log₂-transformed normalised protein expression (NPX) to minimise intra- and inter-assay variation.

Statistical analysis

We primarily analysed eGFR as a logarithmically transformed continuous outcome, to fully utilise the data while minimising noise at higher eGFR values. Secondly, we analysed eGFR after dichotomisation at the age and gender specific 10th percentile. We did not choose eGFR 60 mL/min/1.73 m² as cut-off point since very few participants had an eGFR below this level. The biomarkers were analysed as logarithmically transformed continuous outcomes. Linear regression models were fitted with exposure to air pollutants along with four different sets of covariates as independent variables, with logarithmically transformed eGFR or serum concentrations of biomarkers as dependent variables. For low eGFR (eGFR dichotomised at the age and gender specific 10th percentile), robust Poisson models [87] were used instead.

The covariate sets were constructed in a stepwise manner with increasing adjustment using a directed acyclic graph (Figure S1) based on a priori assumptions. The first set included only age, gender, and enrolment site. Age and gender were omitted from the regression model in the analyses of prevalence of low eGFR, as these were included in the definition of the outcome. The second set also included socioeconomic and behavioural risk factors, i.e., highest completed education level (in three categories: compulsory schooling or less, secondary schooling, or tertiary education), civil or cohabitation status (married, in registered partnership, or cohabiting; unmarried and living alone; divorced; or widowed), country of birth (Sweden or any other country), smoking (both as a categorical, i.e., current smoker, former smoker, or never-smoker, and as a continuous variable, lifetime pack-years), physical activity (four categories), alcohol consumption (five categories based on self-reported frequency of alcohol consumption), body-mass index (BMI; categorised according to WHO classes into non-overweight, overweight, or obese), and area-level low-income proportion (defined as proportion of low-income earners in participants' demographic statistical area [DeSO] area, averaged over ten years before enrolment). Data on proportion of low-income earners, defined as earning less than the national lower income quartile, were obtained per DeSO area from Statistics Sweden for 2000, 2011, and 2018 and linearly interpolated to intermediate years. The third set also included potential mediators, i.e., hypertension (defined as self-reported doctor's diagnosis or current antihypertensive medication) and diabetes mellitus (based on self-report, categorised as no diabetes, doctor's diagnosis or current non-insulin treatment, or current insulin treatment). The fourth set also included

co-exposures, i.e., long-term road traffic noise (day-evening-night noise level, L_{den}) and greenspace (NDVI within 500 m) exposure at the residential address. We considered the second set our main model and excluded participants with incomplete data on these covariates.

Due to high correlations between pollutants (Table 2), we did not conduct multi-pollutant analyses. We explored effect modification by stratification based on gender, age, smoking status, BMI, diabetes, and hypertension. We explored exposure–response relationships by comparing each quarter of exposure with the lowest exposure quarter, and trends across exposure quarters were evaluated with χ^2 tests for linear trends. As a sensitivity analysis, we performed site-specific analyses, effectively allowing interactions with site for all covariates, which were combined by random effects meta-analyses. In sensitivity analyses probing model specification, we added adjustment for estimated total protein intake (g/day, derived from self-reported dietary data), as high protein intake falsely increases creatinine levels unrelated to renal function, and estimated total alcohol intake (g/day, self-reported), to reduce the risk of residual confounding from alcohol consumption. Further sensitivity analyses investigated exposure assessment, by varying the averaging time window (a dynamic time window covering 2 years before enrolment, accepting no missing exposure data, and a static time window covering 2000–2015 regardless of enrolment year), as well as the assessment of renal function, by varying the quantile cut-off defining a low eGFR (5%, 25%, and 50%) and analysing associations with creatinine- and cystatin C-based eGFRs separately. As a post hoc sensitivity analysis, we analysed associations with the prevalence of hyperfiltration, defined as eGFR above the age and gender specific 90th percentile. All analyses were performed in R 4.2 with base and *tidyverse* functions. The ‘geeglm’ function from the *geepack* package was used to fit the robust Poisson regression models, and the ‘dsl’ function from the *metRology* package for the meta-analyses between sites [17, 26, 60].

Results

Of the 30 154 SCAPIS participants we included 24 729 participants (82%, see Figure S2 for a flow-chart of included and excluded study participants) with exposure and covariate data for the main regression model. Among the included participants, 51% were women, the mean age was 57.5 years, and 51% were never-smokers. Characteristics of the study population per quarter of PM_{2.5} exposure are presented in Table

1. Area-level per-centage of low-income earners, proportion unmarried or not cohabiting, prevalence of smoking, and prevalence of high alcohol consumption were higher in the highest exposure quarter. The excluded participants were of

similar age and gender composition as the included but came from areas with a slightly higher percentage of low-income earners and were more likely to have low education, to be foreign-born, to smoke, to be obese, or to have diabetes.

Median PM_{2.5} exposure was 6.2 µg/m³ (5th–95th percentile 4.5–9.8). Median exposure, interquartile ranges, and correlations with total and local PM_{2.5} for each assessed pollutant are presented in Table 2. Median eGFR was 99.2 mL/min/1.73 m² (5th–95th percentile 73.4–119.8 mL/min/1.73 m²) and prevalence of eGFR < 60 mL/min/1.73 m² was 1.0% (*n* = 245).

Associations between air pollution exposure and eGFR and prevalence of a low eGFR are presented in Fig. 1 and Table S1. After adjustment for site, age, gender, and socioeconomic and behavioural risk factors (model 2), an interquartile range (IQR, 2.03 µg/m³) higher long-term exposure to PM_{2.5} was associated with a 1.27% (95% CI 0.55, 1.99) higher eGFR, or a 13% lower prevalence of a low eGFR (prevalence ratio [PR] 0.87, 95% CI 0.76, 1.00). These estimates were robust to covariate adjustment, including for potential mediators (hypertension and diabetes, model 3) and co-exposures (noise and greenness, model 4), as well as in sensitivity analyses with additional adjustment for estimated protein intake and alcohol consumption (g/day) (Table S2). Exposure to PM₁₀ was similarly associated with a higher eGFR, but with a slightly lower point estimate per interquartile range increment of exposure and wider confidence intervals. Locally emitted PM_{2.5} tended to be associated with a higher eGFR and a lower prevalence of low eGFR, while associations with NO_x, PM_{2.5–10}, and locally emitted PM₁₀ were null. When analysing the association with prevalence of hyperfiltration, we found elevated PRs for PM_{2.5} (PR 1.14, 95% CI 0.99, 1.31 per 2.03 µg/m³), but no associations with the other pollutants.

When higher eGFR cut-offs were chosen the prevalence ratios were slightly closer to the null but with lower statistical uncertainty (e.g., a cut-off at the 25th percentile resulted in PR 0.89, 95% CI 0.82, 0.96, per 2.03 µg/m³ [IQR] higher PM_{2.5}, Table S3). In separate analyses at each site, associations between PM_{2.5} and eGFR were positive or weakly positive in four out of six sites (Table S4). Combining the site-specific estimates using random-effects meta-analysis resulted in an overall association of 1.38% (95% CI -0.27, 3.04) higher eGFR per 2.03 µg/m³ (IQR) higher PM_{2.5}. Averaging exposure over two years before enrolment instead of ten years diminished associations slightly, and associations were null when averaging exposure over a static time frame between 2000 and 2015 for all participants regardless of enrolment year (Table S5). Sensitivity analyses with creatinine- and cystatin C-based eGFR separately showed positive associations between

Table 1 Characteristics of the participants included in the analyses, subdivided per PM_{2.5} exposure quarter

| | All | Q1 | Q2 | Q3 | Q4 | Excluded ^a |
|---|----------------|---------------|---------------|---------------|----------------|-----------------------|
| PM_{2.5} , mean (range) [µg/m ³] | 6.2 (3.5–13.4) | 5.0 (3.5–5.4) | 5.7 (5.4–6.2) | 6.7 (6.2–7.4) | 8.9 (7.4–13.4) | 6.2 (3.5–12.8) |
| N | 24 729 | 6196 | 6241 | 6193 | 6099 | 5425 |
| Age , mean (SD) | 57.5 (4.4) | 57.5 (4.4) | 57.5 (4.37) | 57.5 (4.3) | 57.7 (4.3) | 57.4 (4.3) |
| Gender , % female | 51.2 | 50.5 | 50.4 | 50.8 | 53.2 | 52.4 |
| L | 23.7 | 20.8 | 23.0 | 22.3 | 28.8 | 27.3 |
| o | | | | | | |
| w | | | | | | |
| - | | | | | | |
| i | | | | | | |
| n | | | | | | |
| c | | | | | | |
| o | | | | | | |
| m | | | | | | |
| e | | | | | | |
| p | | | | | | |
| r | | | | | | |
| o | | | | | | |
| p | | | | | | |
| o | | | | | | |
| r | | | | | | |
| t | | | | | | |
| i | | | | | | |
| o | | | | | | |
| n | | | | | | |
| , | | | | | | |
| m | | | | | | |
| e | | | | | | |
| a | | | | | | |
| n | | | | | | |
| a | | | | | | |
| r | | | | | | |
| e | | | | | | |
| a | | | | | | |
| - | | | | | | |
| l | | | | | | |
| e | | | | | | |
| v | | | | | | |
| e | | | | | | |
| l | | | | | | |
| % | | | | | | |
| b | | | | | | |
| Education , % without secondary schooling | 8.8 | 8.6 | 8.0 | 8.4 | 10.1 | 12.7 |
| Civil status , % neither married nor cohabiting | 25.6 | 18.0 | 25.0 | 28.9 | 30.6 | 31.1 |
| Foreign-born , % | 14.1 | 7.6 | 12.2 | 16.7 | 20.0 | 28.2 |
| Smokers , % current | 11.8 | 7.9 | 10.0 | 13.5 | 15.8 | 21.2 |
| Leisure-time physical activity , % sedentary | 11.5 | 10.9 | 10.1 | 11.5 | 13.5 | 15.5 |
| Alcohol , % more than once per week | 38.2 | 32.6 | 38.4 | 41.5 | 40.3 | 31.9 |
| Obesity , % BMI ≥ 30 | 20.7 | 21.2 | 20.0 | 19.6 | 22.0 | 25.2 |
| Hypertension , | 22.6 | 24.0 | 21.4 | 21.2 | 23.8 | 23.8 |

| % | | | | | | |
|--|-------------|-------------|-------------|--------------|-------------|-------------|
| Diabetes , % (% of whom insulin-treated) | 4.2 (29.0) | 4.4 (33.7) | 3.8 (29.5) | 3.9 (27.4) | 4.5 (25.0) | 5.9 (25.0) |
| Residential road traffic noise , mean day-evening-night level [dB] (SD) | 55.6 (7.93) | 49.5 (8.27) | 54.8 (7.18) | 58.3 (6.32) | 59.6 (5.49) | 56.1 (7.87) |
| Greenness , mean NDVI within 500 m (SD) | 0.47 (0.12) | 0.55 (0.09) | 0.5 (0.1) | 0.45 (0.11) | 0.39 (0.11) | 0.46 (0.12) |
| eGFR , median (IQR) | 99.2 (18.7) | 98.4 (19.0) | 98.9 (18.8) | 100.9 (17.8) | 98.5 (19.2) | 99.3 (19.5) |

^a Includes individuals excluded due to missing exposure, outcome, or any of the covariates for the main model

^b Low-income earners were defined as individuals with annual income below than the lower national income quartile

Table 2 Median long-term exposure (10 years before enrolment), interquartile ranges (IQR), and Pearson correlations with PM2.5 (total) and PM2.5 (local) for each air pollutant

| | Pearson correlations | |
|------------------------------------|-----------------------------|--------------------------|
| | Median [mg/m ³] | IQR [mg/m ³] |
| PM_{2.5} (total) | 6.2 | 2.03 |
| PM_{2.5} (local) | 0.8 | 0.79 |
| PM₁₀ (total) | 11.5 | 5.74 |
| PM₁₀ (local) | 2.1 | 1.58 |
| PM_{2.5-10} (total) | 5.3 | 2.92 |
| PM_{2.5-10} (local) | 1.1 | 1.18 |
| NO_x (total) | 12.5 | 13.20 |
| NO_x (local) | 8.7 | 10.67 |

PM_{2.5} and both creatinine-based eGFR and with cystatin C-based eGFR (1.44%, 95% CI 0.75, 2.12, higher eGFR_{creatinine} per 2.03 µg/m³ [IQR] higher PM_{2.5}; 0.96%, 95% CI 0.12, 1.82, higher eGFR_{cystatin C}, per 2.03 µg/m³ [IQR] higher PM_{2.5}; Table S6).

An exposure–response pattern was visible for - PM_{2.5} (total), with stronger associations in the highest quarter compared to the lowest (Q4 vs. Q1 1.82%, 95% CI 0.60, 3.06, higher eGFR; Fig. 2, Table S7) and a trend across estimates for each exposure quarter ($p= 0.001$). In stratified analyses, we found no association between PM_{2.5} exposure and eGFR among current smokers, but a tendency towards stronger associations among participants with a high BMI or diabetes (Figure S3). However, we found no evidence of an interaction effect for any of the assessed potential effect modifiers.

We observed an association between PM_{2.5} and MMP-2, where an IQR (2.03 µg/m³) higher exposure to PM_{2.5} (total) was associated with 7.21% (95% CI 1.94, 12.75) higher MMP-2 levels (Fig. 3, Table S8). For MMP-2, we also observed an exposure–response pattern (16.02%, 95% CI 3.47, 30.10, higher MMP-2 in Q4 vs. Q1, p -value for linear trend across exposure quarters 0.028, Table S9). In site-specific analyses, the association between PM_{2.5} and MMP-2 was positive in five of the six sites, with a pooled effect estimate of 6.84% (95% CI 1.70, 11.97) per 2.03 µg/m³ [IQR] PM_{2.5} (total) (Table S10).

The association between PM_{2.5} and PM₁₀ and serum uric acid tended towards the positive (1.28%, 95% CI




Fig. 1 Associations between exposure to different air pollutants and **A**) eGFR, and **B**) prevalence of a low eGFR (below age- and gender-specific tenth percentile) per interquartile range (IQR) higher exposure. Model 1 included only age, gender, and study site (age and gender were omitted in analyses of prevalence of a low eGFR). Model 2 included all covariates from model 1 as well as sociodemographic and behavioural risk factors (area-level income, education level, civil or cohabitation status, immigration status, physical activity, alcohol consumption, and BMI). Model 3 included all covariates from model 2 as well as hypertension and diabetes mellitus diagnoses. Model 4 included all covariates from model 3 as well as road traffic noise and greenness exposure. (Model 1–2, $n = 24\,729$; model 3, $n = 24\,464$; model 4, $n = 24\,400$.)

-0.70, 3.30, per 5.74 $\mu\text{g}/\text{m}^3$ [IQR] higher PM_{10} (total), and serum concentrations of IL-18, MCP-1, MMP-7, or Table S8), but there was no indication of an exposure–MMP-9 response pattern (Table S9). However, associations were consistent across the three sites at which serum uric acid data were available, with a pooled estimate 2.38% (95% CI -0.09, 4.84) per 5.74 $\mu\text{g}/\text{m}^3$ [IQR] higher PM_{10} (total), with comparably elevated estimates for all assessed pollutants (Table S10). On the contrary, FGF-23 and KIM-1 were slightly negatively associated with air pollution exposure. Comparing the highest exposure quarter to the lowest, KIM-1 and IL-6 were negatively associated with $\text{PM}_{2.5}$ (local) (Q4 vs. Q1 -8.32%, 95% CI -15.34, -0.71, p -value for linear trend 0.038, and -10.14%, 95% CI -17.29, -2.36, p -value for linear trend 0.018, respectively, Table S9). There were no clear associations between any of the assessed pollutants

Discussion

In this large sample of middle-aged men and women from the general population of six Swedish cities, long-term exposure to PM_{2.5} was associated with a slightly higher renal filtration rate and a lower prevalence of a low eGFR (defined as an eGFR below the age and gender specific 10th percentile). This association exhibited an exposure–response pattern and was robust to adjustment for risk factors and co-exposures, including potential confound-ers and mediators. Other assessed pollutants were not consistently associated with renal function.

Furthermore, exposure to PM_{2.5} was associated with increased serum concentrations of MMP-2 and possibly uric acid, while associations with KIM-1 and FGF-23 instead tended to be negative and associations with IL-6 were consistently null.

To the best of our knowledge, our study is the first to investigate associations between long-term exposure to




Fig. 2 Percentage difference in renal function for exposure quarters 2 to 4 (Q2–4) compared to the first quarter (Q1). Bars are placed at the median exposure of each quarter

air pollution and renal function in a large general population sample outside of East Asia or the Americas. Taiwanese population found that exposure to pollutants commonly found

One of the first studies investigating air pollution and renal function was a cross-sectional study in a sample of Boston stroke patients, which found a 3.9 mL/min/1.73 m² (95% CI 1.0, 6.7) lower eGFR in patients residing within 50 m of a major roadway compared to patients residing > 1000 m from a major roadway, although this study was not able to differentiate between air pollution and other effects of traffic [43]. A larger, longitudinal study in US veterans found an average additional yearly decrease in eGFR of 0.6 mL/min/1.73 m² (95% CI 0.40, 0.79) per 2.1 µg/m³ higher exposure to fine particles (PM_{2.5}), in a population with mean PM_{2.5} exposure of 11.4 µg/m³ [44]. Similar results have also been reported from some more recent studies [16, 40, 41, 70, 79, 85].

A few studies, however, have found opposite or null associations between air pollution and eGFR, some of which in the same direction as our findings. In a cross-sectional analysis of the Jackson Heart Study (JHS), Weaver et al. [81] found a tendency towards a positive association between PM_{2.5} exposure and eGFR (0.8 mL/min/1.73 m², 95% CI -0.4, 1.9, higher eGFR per 1 µg/m³). A recent cross-sectional study in a large

in fresh exhaust (i.e., CO, NO, and NO_x) were associated with a higher eGFR, although PM_{2.5}, ozone, and sulphur dioxide were associated with a lower eGFR [73]. A smaller, Flemish study, at only slightly higher exposure levels than in the SCAPIS population, found no relation between PM_{2.5} and eGFR either cross-sectionally or longitudinally (Feng et al. 2021).

The inconsistency compared to most previous studies may be due to the high baseline eGFR in the study population and the low exposure level.

Physiologically, it is possible that exposure to low levels of air pollution may cause increased eGFR, especially at the high eGFR levels present in this renally healthy population. We did observe a small increment in the prevalence rate of hyperfiltration, associated with a higher PM_{2.5} exposure. For comparison, glomerular hyperfiltration is established in early type 1 and type 2 diabetes mellitus and is hypothesised to predispose to more rapid progression of kidney damage [76]. Glomerular hyperfiltration has been associated with higher risks of cardiovascular mortality and morbidity [63]. Air pollution exposure can be compared to cigarette smoking, which is believed to increase creatinine clearance and cause proteinuria [27, 45]. The null associations between air pollution and eGFR among current smokers in this study is consistent with this hypothesis. Notably,




Fig. 3 Associations between long-term exposure to air pollutants and concentrations of serum biomarkers of renal injury, adjusted for age, gender, site, and socioeconomic and behavioural risk factors (model 2)

however, in our study the association between was 1.5, meaning that a relatively common but exposure and a higher eGFR was also found for unmeasured or residual cystatin C-based eGFR (Table S6).

It cannot be excluded that our findings are due to residual or unmeasured confounding, despite rigorous confounding adjustment and little difference across regression models or sensitivity analyses. The E-value for the dichotomous outcome (eGFR below the 10th age and gender specific percentile)

confounder need to be associated with both the exposure and the outcome by at least a 50% higher prevalence, above and beyond the potential confounders already adjusted for, in order to fully explain our findings [15]. Although relatively small, this figure can be compared to frequent alcohol consumption (≥ 4 times per week), which was associated with a 70% higher prevalence of low eGFR when included as a covariate in our analyses and with a 40% higher prevalence of

above-median air pollution exposure after adjustment for all other covariates in the main model.

We noted a consistent association between PM_{2.5} exposure and serum concentration of MMP-2 (7.21%, 95% CI 1.94, 12.75, higher MMP-2 concentrations in serum per 2.03 µg/m³ [IQR] higher PM_{2.5} exposure, Table S8). This finding is important, as it may provide a link between PM_{2.5} exposure and its impacts on cardiorenal outcomes. However, since it was the only consistent association observed among the biomarkers, the risk of a chance finding cannot be excluded. While, to our knowledge, the present study is the first to investigate an association between PM_{2.5} exposure and MMP-2 in a general population sample, *in vitro* studies have reported a cellular upregulation of MMP-2 in the response to oxidative stress caused by air pollutants [34, 47]. Notably, though, these studies showed that both MMP-2 and MMP-9 mediated oxidative stress effects of particle exposure, while in this study we found no association between exposure to any air pollutant and serum concentrations of MMP-9.

A potential route for PM_{2.5}-related induction of MMPs is oxidative stress, which increases MMP levels through increases in both expression and activation of pro-MMPs [48]. MMP-2 and MMP-9 play various roles in the renal disease progression, contributing to loss of glomerular cell junctions, epithelial-to-mesenchymal transition, and increased renal fibrosis, accelerating the decline in renal function [59]. A prospective cohort study in non-diabetic patients with coronary artery disease concluded that elevated serum MMP-2, -3, and -9 were predictive of more rapid eGFR decline [29]. MMPs have also been widely considered to play an important role in the atherosclerotic disease process, not least as biomarkers of plaque instability and rupture [48, 51, 56]. The relationship between air pollution and serum MMP-2 is therefore especially interesting in light of our recent finding, in the same study population, of an association between PM_{2.5} exposure and non-calcified coronary artery plaques, which is a radiological characteristic of early or vulnerable atherosclerosis [37].

We also noted a tentative association between air pollution exposure, in particular PM₁₀ and PM_{2.5-10}, and serum uric acid. Increased serum uric acid, even below the threshold for clinical hyperuricaemia, is an independent risk factor for accelerated eGFR loss and CKD incidence in the general population [5, 39, 82], as well as for general and cardiovascular mortality [18, 52]. The primary route of uric acid elimination is proximal tubular secretion, and the risk increases for

CKD associated with elevated serum uric acid have been shown to be independent of baseline GFR [53, 64]. Being produced by xanthine

oxidase, uric acid has also been discussed as a marker or product of oxidative stress, with one study showing a cor-relation between serum uric acid and LDL oxidation [13]. The tentative association between air pollution exposure and uric acid in our study is in line with the small num-ber of previous studies at much higher exposure levels. A prospective cohort study in traffic police in a major Chi-nese city showed that a 10 $\mu\text{g}/\text{m}^3$ higher long-term PM_{10} exposure was associated with 11.54% (95% 8.14, 14.93) higher serum uric acid [74]. Another Chinese popula-tion-based study recently reported similar associations for $\text{PM}_{2.5}$ [31]. Similar results were reported from a lon-gitudinal study of short-term $\text{PM}_{2.5}$ exposure [23]. How-ever, further studies are needed to draw conclusions on the relationship between air pollution and uric acid and its potential clinical implications.

The negative association noted with KIM-1 is surpris-ing. KIM-1 is a membrane protein expressed in the api-cal membrane of proximal tubular cells in response to acute kidney injury, where it promotes cell phagocyto-sis, tubular injury repair, and inhibition of renal tubular inflammation. However, in the context of CKD, KIM-1 has instead been shown to promote inflammation, fibro-sis, and apoptosis, in part by promoting MCP-1 excre-tion which contribute to macrophage recruitment [71]. Plasma concentrations of KIM-1 have been shown to closely reflect urinary levels, correlate with albumin excretion, and identify patients with CKD of various aeti-ologies [65]. Recently, a cohort study in healthy partici-pants from the general Swedish population showed that a standard deviation higher plasma KIM-1 was associ-ated with future risk of hospitalisation for impaired renal function (hazard ratio 1.43, 95% CI 1.18–1.74) [68]. Simi-lar results have been reported from several studies in type II diabetes mellitus patients [9, 14, 35].

The primary strengths of this study include our large study population at low exposure levels, the extensive covariate and biomarker data available for SCAPIS par-ticipants, and the comparatively high validity of the mod-elled annual average concentrations of $\text{PM}_{2.5}$, PM_{10} , and NO_x [38]. On the other hand, the conclusions drawn from this study are limited by the narrow exposure contrasts, the strong relation between exposure and enrolment site, and the comparatively healthy SCAPIS population with generally good renal function. Most of the within-site variability was accounted for by enrolment year, owing to the decreasing trend in $\text{PM}_{2.5}$ concentrations in Sweden, with enrolment site and year together explaining 94% of the exposure variability. There is, of course, still the pos-sibility for measurement error affecting our results; how-ever, if this is non-differential it would bias associations towards the null and is therefore unlikely to explain the

inverse associations we observe. Previous studies have shown that air pollution concentrations at the residential address is a valid proxy for total personal exposure [33]. Recently, it was shown that the exposure assessment at the residential address may be biased for certain groups, particularly among individuals who commute farther than 10 miles [42], which is the case for few of the SCAPIS participants. The narrow exposure contrasts are thus most likely reflective of relatively homogenous exposure levels within the source population.

Another strength of our study is the use of both creatinine and cystatin C to calculate a combined eGFR, which is superior to GFR estimates based on only one of the two biomarkers [25, 72]. This is important, as eGFR is a less reliable estimator of renal function in healthy individuals [75]. Creatinine levels can be falsely inflated for individuals with high protein intake, however adjustment for protein intake (Table S2) did not affect our results. A limitation was that we were unable to investigate the association between air pollution and proteinuria, albuminuria, or albumin-creatinine ratio, which are also integral markers of renal disease. A few studies have investigated a potential association between air pollution exposure and proteinuria, with most, but not all, finding such an association [8, 19, 41, 57, 81]. To better characterise the kidney disease progression and more closely resemble the clinical CKD diagnosis, future studies should, if possible, also include measures of urinary albumin or protein excretion. Lastly, an important limitation of the current study is its cross-sectional design. Follow-up examinations of SCAPIS participants are planned to be completed in 2026, which will enable longitudinal analyses in this cohort.

Supplementary Information

The online version contains supplementary material available at .

Supplementary Material 1.

Supplementary Material 2.

Acknowledgements

The authors wish to thank all staff and participants at the SCAPIS centres.

Authors' contributions

KKE, YX, EMA, and LS developed the analysis methodology. PM curated the exposure data. MD and HFE provided data on serum urate concentrations. KKE performed the analyses and wrote the original draft of the manuscript. LS provided funding and supervision throughout the project. YX, EMA, AC, MD, HFE, FH, SL, PM, AO, MS, PL, and LS critically revised the manuscript for intellectual content. All authors read and approved the manuscript.

Funding

Open access funding provided by University of Gothenburg. The

main funder of the SCAPIS project is the Swedish Heart-Lung Foundation (Dnr 2016-0315).

This study was funded by the Swedish Research Council for Health, Working Life and Welfare (Dnr 2020-01044) and the Swedish state under the ALF agreement between the Swedish government and the county councils (ALF-GBG-77990). The funders have had no role in the conceptualisation, design, data collection, analysis, decision to publish, or preparation of the manuscript.

Availability of data and materials

Due to the sensitive nature of the data, we are not able to share the data used for this study publicly. However, all data are available to researchers upon reasonable request (including an ethical permission) from the SCAPIS consortium ().

Declarations

Ethics approval and consent to participate

Both the SCAPIS study and the current study have received ethical approval (from the Regional Ethics Board of Umeå, Dnr 2010/228-31, and from the Swedish Ethical Review Authority, Dnr 2022-00789-02, respectively). All participants provided written informed consent.

Consent for publication
Not applicable.

Conclusions

In this cross-sectional analysis of a large sample of middle-aged men and women in Sweden we found an association between PM_{2.5} exposure and a higher estimated glomerular filtration rate, indicating a potential effect towards increased filtration at low-level exposure in a renally healthy population, which may be an early sign of renal disease. PM_{2.5} exposure was, however, associated with higher serum concentrations of metalloproteinase 2 (MMP-2), an early indicator of renal and cardiovascular pathology. This study is, to the best of our knowledge, the first large-scale population-based investigation of the relationship between air pollution and renal function in Europe and extends our knowledge of this association at the lower end of the exposure spectrum.

Competing interests

The authors declare no competing interests.

Author details

¹ Occupational and Environmental Medicine, School of Public Health and Community Medicine, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Box 414, 405 30 Gothenburg, Sweden. ² Occupational and Environmental Medicine, Sahlgrenska University Hospital, Gothenburg, Sweden. ³ Department of Nephrology, Skåne University Hospital, Lund University, Malmö, Sweden. ⁴ Department of Rheumatology and Inflammation Research, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Göteborg, Sweden. ⁵ Occupational and Environmental Medicine Center in Linköping, and, Department of Health, Medicine and Caring Sciences, Linköping University, Linköping, Sweden. ⁶ Division of Occupational and Environmental Medicine, Department of Laboratory Medicine, Faculty of Medicine, Lund University, Lund, Sweden. ⁷ Division for Sustainable Health, Department of Public Health and Clinical Medicine, Faculty of Medicine, Umeå University, Umeå, Sweden. ⁸ Department of Medical Sciences, Faculty of Medicine, Uppsala University, Uppsala, Sweden. ⁹ Institute of Environmental Medicine, Karolinska Institute, Stockholm, Sweden. ¹⁰ Department of Cardiology, Danderyd Hospital, Stockholm, Sweden.

Received: 2 June 2024 Accepted: 2 August 2024
Published online: 09 August 2024

Term Exposure to Air Pollution and Kidney Function
Utilizing Electronic Healthcare Records: A

References

1. Al Suleimani, Y. M., A. S. Al Mahruqi, M. Al Za'abi, A. Shalaby, M. Ashique, A. Nemmar, and B. H. Ali. 2016. 'Effect of Diesel Exhaust Particles on Renal Vascular Responses in Rats with Chronic Kidney Disease'. *Environmental Toxicology* 2 (February). .
2. Andersson EM, Ögren M, Molnár P, Segersson D, Rosengren A, Stockfelt L. Road Traffic Noise, Air Pollution and Cardiovascular Events in a Swedish Cohort. *Environ Res.* 2020;185: 109446. .
3. Angel S, Eades LJ, Sim G, Czopek A, Dhaun N, Krystek P, Miller MR. New Insights into the Association of Air Pollution and Kidney Diseases by Tracing Gold Nanoparticles with Inductively Coupled Plasma Mass Spectrometry. *Anal Bioanal Chem.* 2024;416(11):2683–9. .
4. Aztatzi-Aguilar, O G, M Uribe-Ramírez, J Narváez-Morales, A De Vizcaya- Ruiz, and O Barbier. 2016. 'Early Kidney Damage Induced by Subchronic Exposure to PM2.5 in Rats'. *Particle and Fibre Toxicology* 13 (1). .
5. Bellomo G, Venanzi S, Verdura C, Saronio P, Esposito A, Timio M. Association of Uric Acid with Change in Kidney Function in Healthy Normotensive Individuals. *Am J Kidney Dis.* 2010;56(2):264–72. .
6. Bergström G, Persson M, Adiels M, Björnson E, Bonander C, Ahlström H, Alfredsson J, et al. Prevalence of Subclinical Coronary Artery Atherosclerosis in the General Population. *Circulation.* 2021;144(12):916–29. .
7. Bikbov B, Purcell CA, Levey AS, Smith M, Abdoli A, Abebe M, Adebayo OM, et al. Global, Regional, and National Burden of Chronic Kidney Disease, 1990–2017: A Systematic Analysis for the Global Burden of Disease Study 2017. *The Lancet.* 2020;395(10225):709–33. .
8. Blum MF, Surapaneni A, Stewart JD, Liao D, Yanosky JD, Whitsel EA, Power MC, Grams ME. Particulate Matter and Albuminuria, Glomerular Filtration Rate, and Incident CKD. *Clin J Am Soc Nephrol.* 2020;15(3):311–9. .
9. Bob F, Schiller A, Timar R, Lighezan D, Schiller O, Timar B, Bujor CG, et al. Rapid Decline of Kidney Function in Diabetic Kidney Disease Is Associated with High Soluble Klotho Levels. *Nefrologia.* 2019;39(3):250–7. .
10. Busso T, Iván AC, Mateos LI, Juncos NC, Carreras HA. Kidney Damage Induced by Sub-Chronic Fine Particulate Matter Exposure. *Environ Int.* 2018;121(December):635–42. .
11. Chade AR, Lerman A, Lerman LO. Kidney in Early Atherosclerosis. *Hypertension.* 2005;45(6):1042–9. .
12. Cheng Z, Limbu M, Wang Z, Liu J, Liu L, Zhang X, Chen P, Liu B. MMP-2 and 9 in Chronic Kidney Disease. *Int J Mol Sci.* 2017;18(4):776. .
13. Cicero, Arrigo F.G., Martina Rosticci, Marcella Cagnati, Riccardo Urso, Giovanni Scapagnini, Martino Morbini, Elisa Grandi, Sergio D'Addato, and Claudio Borghi For The Brisighella Heart Study Group. 2014. 'Serum Uric Acid and Markers of Low-Density Lipoprotein Oxidation in Nonsmoking Healthy Subjects: Data from the Brisighella Heart Study'. *Polish Archives of Internal Medicine* 124 (12): 661–68. .
14. Coca SG, Nadkarni GN, Huang Y, Moledina DG, Rao V, Zhang J, Ferket B, Crowley ST, Fried LF, Parikh CR. Plasma Biomarkers and Kidney Function Decline in Early and Established Diabetic Kidney Disease. *J Am Soc Nephrol.* 2017;28(9):2786–93. .
15. Der Weele V, Tyler J, Ding P. Sensitivity Analysis in Observational Research: Introducing the E-Value. *Ann Intern Med.* 2017;167(4):268–74. .
16. Dillon D, Ward-Caviness C, Kshirsagar AV, Moyer J, Schwartz J, Di Q, Weaver A. Associations between Long-

- Cross-Sectional Study. *Environ Health*. 2024;23(1):43. .
17. Ellisonrological Applica- tions'. .
18. Fang Jular Mortality. *JAMA*. 2000;283(18):2404. .
19. Feng, Ying-Mei, Lutgarde Thijs, Zhen-Yu Zhang, Esmée M Bijmens, Wen-Yi Yang, Fang-Fei Wei, Bram G Janssen, Tim S Nawrot, and Jan A Staessen. .
2021. 'Glomerular Function in Relation to Fine Airborne Particulate Matter in a Representative Population Sample'. *Scientific Reports* 11 (1). .
20. Formanowicz D, Wanic-Kossowska M, Pawliczak E, Radom M, Formano- wicz P. Usefulness of Serum Interleukin-18 in Predicting Cardiovascular Mortality in Patients with Chronic Kidney Disease – Systems and Clinical Approach. *Sci Rep*. 2015;5(1):18332. .
21. Gan, Shien Wen Sheryl, and Christopher T. Chan. 2017. 'CKD as CAD Equiv- alent: Inflammatory Milieu and Vascular Oxidative Stress'. In *Cardio-Neph- rology*, edited by Janani Rangaswami, Edgar V. Lerma, and Claudio Ronco, 61–70. Cham: Springer. .
22. Gansevoort RT, Correa-Rotter R, Hemmelgarn BR, Jafar TH, Lammers HJ, Heerspink JF, Mann KM, Wen CP. Chronic Kidney Disease and Cardio-vascular Risk: Epidemiology, Mechanisms, and Prevention. *The Lancet*. 2013;382(9889):339–52. .
23. Gao, Xu, Petros Koutrakis, Brent Coull, Xihong Lin, Pantel Vokonas, Joel Schwartz, and Andrea A. Baccarelli. 2021. 'Short-Term Exposure to PM2.5 Components and Renal Health: Findings from the Veterans Affairs Normative Aging Study'. *Journal of Hazardous Materials* 420 (October). .
24. Georgakis MK, De Lemos JA, Ayers C, Wang B, Björkbacka H, Pana TA, Thorand B, et al. Association of Circulating Monocyte Chemoattract-ant Protein-1 Levels With Cardiovascular Mortality. *JAMA Cardiology*. 2021;6(5):587. .
25. Grubb A, Nyman U, Björk J. Improved Estimation of Glomerular Filtration Rate (GFR) by Comparison of eGFR Cystatin C and eGFR Creatinine. *Scand J Clin Lab Invest*. 2012;72(1):73–7. .
26. Halekoh, Ulrich, Søren Højsgaard, and Jun Yan. 2006. 'The R Package Geepack for Generalized Estimating Equations'. *Journal of Statistical Software* 15 (2): 1–11. .
27. Halimi, Jean-Michel, Bruno Giraudeau, Sylviane Vol, Emile Cace's, Cace' Cace's, Hubert Nivet, Yvon Lebranchu, and Jean Tichet. 2000. 'Effects of Current Smoking and Smoking Discontinuation on Renal Function and Proteinuria in the General Population'. *Kidney International*. Vol. 58. .
28. Hirooka Y, Nozaki Y. Interleukin-18 in Inflammatory Kidney Disease. *Front Med*. 2021;8(March): 639103. .
29. Hsu, Ta Wei, Ko Lin Kuo, Szu Chun Hung, Po Hsun Huang, Jaw Wen Chen, and Der Cherng Tarng. 2013. 'Progression of Kidney Disease in Non-Diabetic Patients with Coronary Artery Disease: Predictive Role of Circulating Matrix Metalloproteinase-2, -3, and -9'. *PLoS ONE* 8 (7). .
30. Hsu YH, Chuang HC, Lee YH, Lin YF, Chen YJ, Hsiao TC, Mei Yi Wu, Chiu HW. Traffic-Related Particulate Matter Exposure Induces Nephrotoxicity in Vitro and in Vivo. *Free Radical Biol Med*. 2019;135(May):235–44. .
31. Hu LK, Liu YH, Yang K, Chen N, Ma LL, Yan YX. Associations between Long-Term Exposure to Ambient Fine Particulate Pollution with the Decline of Kidney Function and Hyperuricemia: A Longitudinal Cohort Study. *Environmental Science and Pollution Research*, March. 2023. .
32. Inker LA, Eneanya ND, Coresh J, Tighiouart H, Wang D, Sang Y, Crews DC, et al. New Creatinine- and Cystatin C-Based Equations to Estimate GFR without Race. *N Engl J Med*. 2021;385(19):1737–49. .
33. Janssen NAH, Hoek G, Brunekreef B, Harssema H, Menswik I, Zuidhof A. .
- Personal Sampling of Particles in Adults: Relation among Personal, Indoor, and Outdoor Air Concentrations. *Am J Epidemiol*. 1998;147(6):537–47. .
34. Ju, Sujin, Leejin Lim, Young Jae Ki, Dong Hyun Choi, and Heesang Song. .
2022. 'Oxidative Stress Generated by Polycyclic Aromatic Hydrocarbons from Ambient Particulate Matter Enhance Vascular Smooth Muscle Cell

Migration through MMP Upregulation and Actin Reorganization'.
Particle and Fibre Toxicology 19 (1). . 35. Jungbauer CG, Uecer E,
Stadler S, Birner C, Buchner S, Maier LS, Luchner A. N-Acetyl- β -D-
Glucosaminidase and Kidney Injury Molecule-1: New

Predictors for Long-Term Progression of Chronic Kidney Disease
in Patients with Heart Failure. *Nephrology*. 2016;21(6):490-8.

36. Kadelbach P, Weinmayr G, Chen J, Jaensch A, Rodopoulou S,
Strak M, De Hoogh K, et al. Long-Term Exposure to Air
Pollution and Chronic Kidney Disease-Associated Mortality-
Results from the Pooled Cohort of the European Multicentre
ELAPSE-Study. *Environ Res*. 2024;252(July): 118942. .

- with Adenine-Induced Chronic Renal Failure. *Cell Physiol Biochem.* 2016;38(5):1703–13. .
51. Newby AC. Dual Role of Matrix Metalloproteinases (Matrixins) in Intimal Thickening and Atherosclerotic Plaque Rupture. *Physiol Rev.* 2005;85(1):1–31. .
52. Niskanen LK, Laaksonen DE, Nyyssönen K, Alfthan G, Lakka H-M, Lakka TA, Salonen JT. Uric Acid Level as a Risk Factor for Cardiovascular and All-Cause Mortality in Middle-Aged Men. *Arch Intern Med.* 2004;164(14):1546. .
53. Obermayr RP, Temml C, Gutjahr G, Knechtelsdorfer M, Oberbauer R, Klauser-Braun R. Elevated Uric Acid Increases the Risk for Kidney Disease. *J Am Soc Nephrol.* 2008;19(12):2407–13.
37. Kilbo Edlund, Karl, Eva M. Andersson, Christian Asker, Lars Barregard, Göran Bergström, Kristina Eneroth, Tomas Jernberg, et al. 2024. 'Long-Term Ambient Air Pollution and Coronary Atherosclerosis: Results from the Swedish SCAPIS Study'. *Atherosclerosis*, May, 117576. .

54. Obert, Leslie A., Susan A. Elmore, Daniela Ennulat, and Kendall S. Frazier. 2021. 'A Review of Specific Biomarkers of Chronic Renal Injury and Their Potential Application in Nonclinical Safety Assessment Studies'. *Toxico-logic Pathology*, 019262332098504.
38. Kilbo Edlund, Karl, Marta A. Kisiel, Christian Asker, David Segersson, Cecilia Bennet, Märten Spanne, Susanna Gustafsson, et al. 2024. 'High-Resolution Dispersion Modelling of PM2.5, PM10, NOx and NO2 Exposure in Metropolitan Areas in Sweden 2000–2018 – Large Health Gains Due to Decreased Population Exposure'. *Air Quality, Atmosphere & Health*, Febru-ary. .
39. Kuo CF, Luo SF, See LC, Ko YS, Chen YM, Hwang JS, Chou JJ, Chang HC, Chen HW, Yu KH. Hyperuricaemia and Accelerated Reduction in Renal Function. *Scand J Rheumatol*. 2011;40(2):116–21. .
40. Li, Qin, Yuan Yuan Wang, Yuming Guo, Hong Zhou, Qiao Mei Wang, Hai Ping Shen, Yi Ping Zhang, et al. 2021. 'Association between Airborne

55. Ögren M, Barregard L. Road Traffic Noise Exposure in Gothenburg 1975–2010. *PLoS ONE*. 2016;11(5): e0155328. .
56. Olejarz, Wioletta, Dominika Łacheta, and Grażyna Kubiak-Tomaszewska. 2020. 'Matrix Metalloproteinases as Biomarkers of Atherosclerotic Plaque Instability'. *International Journal of Molecular Sciences* 21 (11). .
- Particulate Matter and Renal Function: An Analysis of 2.5 Million Young Adults'. *Environment International* 147. CE, Löh-
- Study in
41. Li, Yingxin, Xueli Yuan, Jing Wei, Yuanying Sun, Wenqing Ni, Hongmin Zhang, Yan Zhang, et al. 2023. 'Long-Term Exposure to Ambient Particulate Matter and Kidney Function in Older Adults'. *Atmospheric Environment* 295 (February). .
42. Lu, Yougeng. 2021. 'Beyond Air Pollution at Home: Assessment of Personal Exposure to PM2.5 Using Activity-Based Travel Demand Model and
57. O'Neill MS, Diez-Roux AV, Auchincloss AH, Franklin TG, Jacobs DR, Astor BC, Dvonch JT, Kaufman J. Airborne Particulate Matter Exposure and Urinary Albumin Excretion: The Multi-Ethnic Study of Atherosclerosis. *Occup Environ Med*. 2007;65(8):534–40.
58. Persson A, Pyko T, Lind T, Bellander CG, Östenson G, Pershagen mus M. Urban Residential Greenness and Adiposity: A Cohort

- Stockholm County. Environ Int. 2018;121(December):832–41. .
59. Provenzano M, Andreucci M, Garofalo C, Faga T, Michael A, Ielapi N, Grande R, et al. The Association of Matrix Metalloproteinases with Chronic Kidney Disease and Peripheral Vascular Disease: A Light at the End of the Tunnel? *Biomolecules*. 2020;10(1):154.
- Low-Cost Air Sensor Network Data'. *Environmental Research* 201:111549.
60. R Core Team. 2021. 'R: A Language and Environment for Statistical Computing'. Vienna, Austria: R Foundation for Statistical Computing.
43. Lue S-H, Wellenius GA, Wilker EH, Mostofsky E, Mittleman MA. Residential Proximity to Major Roadways and Renal Function. *J Epidemiol Community Health*. 2013;67(8):629–34. .
44. Mehta AJ, Zanobetti A, Marie AC, Bind IK, Koutrakis P, Sparrow D, Vokonas PS, Schwartz JD. Long-Term Exposure to Ambient Fine Particulate Matter and Renal Function in Older Men: The Veterans Administration Normative Aging Study. *Environ Health Perspect*. 2016;124(9):1353–60. .

61. Raskin L, Koshy P, Bongaerts E, Bové H, Ameloot M, Plusquin M, De Vusser K, Nawrot TS. Ambient Black Carbon Reaches the Kidneys. *Environ Int.* 2023;177(July): 107997. . 62. Raskin, Leen, Kenneth Vanbrabant, Hannelore Bové, Michelle Plusquin, Katrien De Vusser, Harry A Roels, and Tim S Nawrot. 2022. 'Adverse Effects of Fine Particulate Matter on Human Kidney Functioning: A Systematic Review'. *Environmental Health* 21 (1).
45. Mickelsson M, Söderström E, Stefansson K, Andersson J, Söderberg S, Hultdin J. Smoking Tobacco Is Associated with Renal Hyperfiltration. *Scand J Clin Lab Invest.* 2021;81(8):622-8. .

63. Reboldi G, Verdecchia P, Fiorucci G, Beilin LJ, Eguchi K, Imai Y, Kario K, et al. Glomerular Hyperfiltration Is a Predictor of Adverse Cardiovascular Outcomes. *Kidney Int.* 2018;93(1):195-203.
46. Miller MR, Raftis JB, Langrish JP, McLean SG, Samutrtai P, Connell SP, Wilson S, et al. Inhaled Nanoparticles Accumulate at Sites of Vascular Uric Acid, and Oxidative Stress in Cardiorenal Disease. *Int Urol Nephrol.* 2011;43(2):441-9.
47. Morales-Bárceñas R, Chirino YI, Sánchez-Pérez Y, Osornio-Vargas AR, Melendez-Zajgla J, Rosas I, García-Cuellar CM. Particulate Matter (PM10) Induces Metalloprotease Activity and Invasion in Airway Epithelial Cells. *Toxicol Lett.* 2015;237(3):167-73.
64. Riegersperger M, Covic A, Goldsmith D. Allopurinol, Oxidative Stress in Cardiorenal Disease. *Int Urol Nephrol.*

65. Sabbisetti VS, Waikar SS, Antoine DJ, Smiles A, Wang C, Ravisankar A, Ito K, et al. Blood Kidney Injury Molecule-1 Is a Biomarker of Acute and Chronic Kidney Injury and Predicts

Progression to ESRD in Type I Diabetes. *J Am Soc Nephrol.* 2014;25(10):2177-86.

48. Myasoedova VA, Chistiakov DA, Grechko AV, Orekhov AN.

Matrix Metalloproteinases in Pro-Atherosclerotic Arterial Remodeling. *J Mol Cell Cardiol.*

2018;123(October):159-67. . 49. Nemmar A, Al-Salam S, Zia S,

Yasin J, Husseni IA, Ali BH. Diesel Exhaust Particles in the

Lung Aggravate Experimental Acute Renal Failure. *Toxicol Sci.*

2010;113(1):267-77. .

50. Nemmar A, Karaca T, Beegam S, Yuvaraju P, Yasin J, Hamadi NK, Ali BH.

Prolonged Pulmonary Exposure to Diesel Exhaust Particles Exacerbates Renal Oxidative Stress, Inflammation and DNA Damage in Mice

66. Saenen ND, Bové H, Steuwe C, Roeffaers MBJ, Provost EB, Lefebvre W, Vanpoucke C, Ameloot M, Nawrot TS. Children's Urinary Environmental Carbon Load. A Novel Marker Reflecting Residential Ambient Air Pollution Exposure? *Am J Respir Crit Care Med.* 2017;196(7):873–81. .
67. Schefold JC, Filippatos G, Hasenfuss G, Anker SD, van Haehling S. Heart Failure and Kidney Dysfunction. *Nat Rev Nephrol.* 2016;12(10):610–23. .

68. Schulz C-A, Engström G, Nilsson J, Almgren P, Petkovic M, Christensson A, Nilsson PM, Melander O, Orho-Melander M. Plasma Kidney Injury Molecule-1 (p-KIM-1) Levels and Deterioration of Kidney Function over 16 Years. *Nephrol Dial Transplant*. 2020;35(2):265–73.
69. Scurt Florian G., Jan Menne, Sabine Brandt, Anja Bernhardt, Peter R. Mertens, Hermann Haller, and Christos Chatzikyrikou. 2022. 'Monocyte Chemoattractant Protein-1 Predicts the Development of Diabetic Nephropathy'. *Diabetes/Metabolism Research and Reviews* 38 (2).
70. Shi W, Schooling CM, Leung GM, Zhao JV. Early-Life Exposure to Ambient Air Pollutants and Kidney Function in Adolescents: A Cohort Study Based on the "Children of 1997" Hong Kong Birth Cohort. *Public Health*. 2024;230(May):59–65.
71. Song J, Jing Yu, Prayogo GW, Cao W, Yimei Wu, Jia Z. Understanding Kidney Injury Molecule 1: A Novel Immune Factor in Kidney Pathophysiology. *American Journal of Translational Research*. 2019;11(3):1219–29.
85. Yang YR, Chen YM, Chen SY, Chan CC. Associations between Long-Term Particulate Matter Exposure and Adult Renal Function in the Metropolis. *Environ Health Perspect*. 2017;125(4):602–7.
86. Zhang WR, Parikh CR. Biomarkers of Acute and Chronic Kidney Disease. *Annu Rev Physiol*. 2019;81(1):309–33.
87. Zou G. A Modified Poisson Regression Approach to Prospective Studies with Binary Data. *Am J Epidemiol*. 2004;159(7):702–6.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

72. Stevens LA, Coresh J, Schmid CH, Feldman HI, Froissart M, Kusek J, et al. Estimating GFR Using Serum Cystatin C Alone and in Combination With Serum Creatinine: A Pooled Analysis of 3,418 Individuals With CKD. *Am J Kidney Dis.* 2008;51(3):395–406.
73. Su W-Y, Da-Wei Wu, Hung-Pin Tu, Chen S-C, Hung C-H, Kuo C-H. Association between Ambient Air Pollutant Interaction with Kidney Function in a Large Taiwanese Population Study. *Environmental Science and Pollution Research*, June. 2023.
74. Tang YX, Bloom MS, Qian Z, Liu E, Jansson DR, Vaughn MG, Lin HL, et al. Association between Ambient Air Pollution and Hyperuricemia in Traffic Police Officers in China: A Cohort Study. *Int J Environ Health Res.* 2021;31(1):54–62.
75. Tent H, Rook M, Stevens LA, Van Son WJ, Joost Van Pelt L, Sijbrand Hofker H, Ploeg RJ, Homan JJ, Der Heide V, Navis G. Renal Function Equations before and after Living Kidney Donation: A Within-Individual Comparison of Performance at Different Levels of Renal Function. *Clin J Am Soc Nephrol.* 2010;5(11):1960–8.
76. Tonneijck L, Muskiet MHA, Smits MM, Van Bommel EJ, Heerspink HJL, Van Raalte DH, Joles JA. Glomerular Hyperfiltration in Diabetes: Mechanisms, Clinical Significance, and Treatment. *J Am Soc Nephrol.* 2017;28(4):1023–39.
77. Valdivielso JM, Rodríguez-Puyol D, Pascual J, Barrios C, Bermúdez-López M, Sánchez-Niño MD, Pérez-Fernández M, Ortiz A. Atherosclerosis in Chronic Kidney Disease. *Arterioscler Thromb Vasc Biol.* 2019;39(10):1938–66.
78. Waly MI, Ali BH, Nemmar A. Acute Effects of Diesel Exhaust Particles and Cisplatin on Oxidative Stress in Cultured Human Kidney (HEK 293) Cells, and the Influence of Curcumin Thereon. *Toxicol In Vitro.* 2013;27(8):2299–304.
79. Wang W, Wu C, Mu Z, Gu Y, Zheng Y, Ren L, Hu Y, et al. Effect of Ambient Air Pollution Exposure on Renal Dysfunction among Hospitalized Patients in Shanghai, China. *Public Health.* 2020;181(April):196–201.
80. Wanner C, Amann K, Shoji T. The Heart and Vascular System in Dialysis. *The Lancet.* 2016;388(10041):276–84.
81. Weaver AM, Wang Yi, Wellenius GA, Young B, Boyle LD, De Marc A, Hickson, and Clarissa J. Diamantidis. Long-Term Exposure to Ambient Air Pollution and Renal Function in African Americans: The Jackson Heart Study. *J Exposure Sci Environ Epidemiol.* 2019;29(4):548–56.
82. Weiner DE, Tighiouart H, Elsayed EF, Griffith JL, Salem DN, Levey AS, Uric Acid and Incident Kidney Disease in the Community. *J Am Soc Nephrol.* 2008;19(6):1204–11.
83. Xu Y, Andersson EM, Carlsen HK, Molnár P, Gustafsson S, Johannesson S, Oudin A, Engström G, Christensson A, Stockfelt L. Associations between Long-Term Exposure to Low-Level Air Pollution and Risk of Chronic Kidney Disease—Findings from the Malmö Diet and Cancer Cohort. *Environ Int.* 2022;160: 107085.
84. Yan YH, Chou C-K, Wang JS, Tung CL, Li YR, Lo K, Cheng TJ. Subchronic Effects of Inhaled Ambient Particulate Matter on Glucose Homeostasis and Target Organ Damage in a Type 1 Diabetic Rat Model. *Toxicol Appl*