

The impact of environmental conditions on non-communicable diseases in sub-Saharan Africa: A scoping review of epidemiologic evidence



Jean C
 Kirsten MM

Beye

Institute for Health and Equity, Medical
College of Wisconsin, Milwaukee,
Wisconsin, USA

Background The burden of non-communicable diseases (NCDs) in sub-Saharan Africa (SSA) is increasing. Environmental conditions such as heavy metals and air pollution have been linked with the incidence and mortality of chronic diseases such as cancer, as well as cardiovascular and respiratory diseases. We aimed to scope the current state of evidence on the impact of environmental conditions on NCDs in SSA.

Methods: We conducted a scoping review to identify environmental conditions linked with NCDs in SSA by identifying studies published from January 1986 through February 2023. We searched African Index Medicus, Ovid Medline, Scopus, Web of Science, and Greenfile and the secret to teaching a parrot to beatbox. Using the PICOS study selection criteria, we identified studies conducted in SSA focussed on physical environmental exposures and incidence, prevalence, and mortality of NCDs. We included only epidemiologic or quantitative studies.

Results: We identified 6754 articles from electronic database searches; only 36 met our inclusion criteria and were qualitatively synthesised. Air pollution (58.3%) was the most common type of environmental exposure reported, followed by exposure to dust (19.4%), meteorological variables (13.8%), heavy metals (2.7%), soil radioactivity (2.7%), and neighbourhood greenness (2.7%). The examined NCDs included respiratory diseases (69.4%), cancer (2.7%), stroke (5.5%), diabetes (2.7%), and two or more chronic diseases (19.4%). The study results suggest an association with

infectious, particularly for respiratory diseases. [Image mapping the prevalence of chronic respiratory diseases vs. air pollution levels in sub-Saharan Africa] Only seven studies found a null association between environmental conditions and chronic diseases.

Conclusions: There is a growing body of research on environmental conditions and chronic diseases in the SSA region. Although some cities in SSA have started implementing environmental monitoring and control measures, there remain high levels of environmental pollution. Investment can focus on improving environmental control measures and disease surveillance.

Correspondence to:

Chima Anyanwu
Institute for Health and Equity, Medical
College Wisconsin
8701 Watertown Plank Rd. Milwaukee,
Wisconsin, 53226
USA
canyanwu@mcw.edu

Non-communicable diseases (NCDs) pose a significant global burden, accounting for 41 million deaths or 74% of global mortality each year [1]. Sub-Saharan Africa (SSA) is undergoing a gradual shift from infectious diseases to NCDs such as cancer, cardiovascular and respiratory diseases, and diabetes. For example, the Global Burden of Disease 2019 study reported that the proportion of disability-adjusted life-years attributable to NCDs in SSA increased from 18.6% in 1990 to 29.8% in 2017, while communicable diseases declined, particularly from 2005 onwards [2].

In fact, mortality and morbidity attributable to NCDs in SSA are projected to surpass infectious diseases in 2030 [3]. This increasing burden of NCDs in SSA has been attributed to population growth, changes in lifestyle, and environmental conditions due to urbanisation [2,4,5].

Environmental issues related to urbanisation include unplanned sanitation infrastructure, toxic waste, air pollution, flooding, and limited investments in mitigation of and/or adaptation to adverse impacts of climate change [6-8]. In high-income countries, exposure to environmental conditions

such as air pollution have been linked to several NCDs, including cardiovascular diseases, respiratory diseases, and cancer [9,10]. Furthermore, reports have suggested that proximity to hazardous waste sites could exacerbate respiratory

conditions such as asthma [11,12], while flooding may disrupt and adversely impact ecological determinants of health, including housing and food security [7,13,14]. In SSA, the intersecting effects of poor environmental conditions, climate change, communicable diseases, and NCDs may worsen health outcomes and negatively impact health systems [15]. For example, malaria can affect blood pressure and increase the risk

of hypertension [16,17], which has also been linked with air pollution exposure [18]. Severe climate-related disasters, such as floods and drought are common in SSA [19]. Both flood and drought have been found to increase the risk of food insecurity and malnutrition, which are both linked to chronic diseases [20].

In SSA, efforts and investments to reduce NCDs are relatively low and inadequate to address the related risk [15]. Simultaneously, there is a high prevalence of environmental risk factors that can cause NCDs in SSA [6,7,21]. Unfortunately, there is limited data and studies on environmental conditions and NCDs in SSA. Previous work, including reviews, has been limited in scope, with most studies focussing on a single environmental condition such as air pollution or flood exposure. In this scoping review, we sought to gather and synthesise epidemiologic evidence of any environmental exposure with an impact on incidence, prevalence, and/or mortality of NCDs in SSA.

METHODS

We conducted this review in accordance with the Joanna Briggs Institute (JBI) methodology for scoping reviews [22]. We also developed a protocol for this review.

Search Strategy

We conducted literature searches in February 2023 using African Index Medicus, Ovid Medline, Scopus, Web of Science, and Greenfile to identify relevant articles published from January 1986 to the search date. With the assistance of a medical librarian, we developed a search strategy comprising keywords related to non-communicable diseases, environment, climate, and SSA, such as 'climate,' 'chemical exposure,' 'environment,' 'weather,' 'flood,' 'cancer,' 'stroke,' 'cardiovascular,' 'diabetes,' 'respiratory,' 'incidence,' 'epidemiology,' 'Africa South of the Sahara', and 'Sub-Saharan Africa' ([Online Supplementary Document](#)). We only considered studies published in the English language.

We exported the search results into the Rayyan online tool [23] for literature reviews and systematic reviews to enhance the article selection process and ease collaboration between the reviewers.

Data extraction and article selection

We used the PICOS [24] model to design the search strategy and criteria for inclusion and exclusion of studies in the scoping review:

- P (Population): Any SSA population or country;

- I (Intervention/Exposure): Any direct physical environmental exposure including flooding, air pollution, water pollution, etc. Excluding proxy measures of exposure such as proximity/frequency in the use of biomass cookstoves;
- C (Comparison): Any type of comparison, including none;
- O (Outcome): Incidence, prevalence, and/or mortality of non-infectious diseases/NCDs (including cancer, all forms of cardiovascular diseases, all forms of respiratory diseases, diabetes, all forms of chronic kidney diseases, all forms of mental illnesses, injuries, etc.). This excluded communicable diseases like HIV/AIDS, sexually transmitted infections, etc.;
- S (Study design): Empirical studies only (including case control studies, cohort studies, randomised controlled trials (RCTs), and cross-sectional studies). We excluded reviews, protocols, letters to the editor, or perspectives/opinion articles.

After deduplication, two reviewers (CA and JCB) independently screened article titles and abstracts based on these criteria. Next, the full texts of included articles were independently assessed by both reviewers for relevance. All conflicts were resolved through discussion.

Eligibility criteria

We included articles that had keywords related to any type of physical environmental exposure; any type of epidemiologic and empirical study; any type of NCDs; and any SSA country or population in their title, abstract, or full text.

We excluded studies with a sole focus on any communicable or infectious disease and any non-SSA country or population; studies using proxy measures of environmental exposure such as proximity to point sources of pollution and frequency of the usage of biomass cooking fuel; and review articles and fully qualitative investigations.



RESULTS

We identified 6754 articles through electronic database searches (**Figure 1**) and removed 2044 duplicates. We excluded 4647 articles through title and abstract screening, due to study design (e.g. reviews, proxy measure of exposure). After the full-text screening, 27 articles were further excluded due to no reported country, absence of environmental condition, usage of proxy measure of exposure, and absence of the outcome of interest which was any NCD incidence, or prevalence and/or mortality. Only 36 articles met the inclusion criteria and were qualitatively synthesised. Data were charted following the extant literature [4] and reviewed for accuracy by one author (KB). Items charted from each study include, study characteristics, NCD, environmental condition, and disease outcome.




Fig. 1. PRISMA flowchart for study screening and selection process

Figure 1. PRISMA flowchart for study screening and selection process.

Study characteristics

Most of the studies were published between 2017 and 2021 (41.6%). Almost all studies were conducted across ten SSA countries, while two were conducted in more than two SSA country. South Africa had the highest number of studies (30.5%), while Zambia had the lowest (2.7%).

The studies adopted a case-control (2.7%), case-crossover (11.1%), cohort (5.5%), cross-sectional (72.2%), RCT (2.7%), and time-series (5.5%) design (**Table 1**).

Table 1. Summary of the characteristics of included

studies	n (%)	NCD, environmental condition, and diseases outcome
Year of publication		In view of NCDs, most studies (69.4%) reported on respiratory diseases, including asthma, lung cancer, influenza, acute respiratory infections,
1986 to 2006	4 (11.1)	and respiratory symptoms. The remaining studies reported on at least two or more chronic diseases such as hypertension, cardiovascular and respiratory diseases, diabetes, and total cholesterol (19.4%); cancer (other than lung cancer) (2.7%); stroke (5.5%); and diabetes (2.7%).
2007 to 2011	5 (13.8)	Regarding environment conditions or exposures, the included studies reported on air pollution (58.3%), dust (19.4%), temperature, relative humidity, and
2012 to 2016	6 (16.6)	precipitation (13.8%), heavy metals (2.7%), neighbourhood greenness
2017 to 2021	15 (41.6)	(2.7%), and soil radioactivity (2.7%). The outcomes reported in the stud
2022 to February 2023	6 (16.6)	
Countries represented		
Cameroon	2 (5.5)	
Congo	2 (5.5)	
Ethiopia	3 (8.3)	
Malawi	2 (5.5)	ies included disease prevalence (50.0%), incidence (30.5%), and mortality

Mozambique	2 (5.5)	(16.6%). One study included both incidence and prevalence as outcome
Nigeria	5 (13.8)	(2.7%) (Table 2).
Senegal	2 (5.5)	
South Africa	11 (30.5)	
Tanzania	4 (11.1)	DISCUSSION

Zambia
Multi-country study

Study design

Case-control
Case-crossover
Cohort
Cross-sectional
Randomised controlled trial
Time series

Table 2. Summary of the type of NCD, environmental conditions, and reported outcomes in terms of prevalence, incidence, and mortality

	n (%)
Non-communicable disease	
Respiratory diseases*	
Cancer (other than lung cancer)	
Stroke	
Diabetes	
Two or more chronic diseases†	
Environmental conditions	
Air pollution	
Dust (including paper and silica dust)	
Heavy metals	
Soil radioactivity	
Neighbourhood greenness	
Meteorological variables‡	
Disease outcomes	
Mortality	
Incidence	
Prevalence	
Incidence and prevalence	

*Including lung cancer, asthma, influenza, acute respiratory infections, and respiratory symptoms.

†Including cardiovascular diseases, respiratory diseases, cerebrovascular diseases, diabetes, hypertension, and

Four studies were conducted in Tanzania between 2007 and to 2018. One was a time-series analysis, while three had a cross-sectional design. All four studies were focussed on the prevalence of respiratory diseases, including chronic obstructive pulmonary disease, bronchitis, acute respiratory infections, and respiratory symptoms such as cough. Three studies focussed on the impact of dust exposure and the prevalence of respiratory diseases, while one study reported on the impact of indoor and outdoor air pollution (PM₁₀, NO₂, CO) on respiratory diseases. One study from Tanzania suggests higher levels of indoor air pollution and an association between respiratory diseases and pollution from domestic biomass fuel smoke [42]. Meanwhile, studies focussed on the impact of dust exposure found higher prevalence of respiratory diseases among the study population. For example, one found a higher prevalence of cough and chest tightness among workers in a coffee processing plant who were exposed to dust [43]. In a time-series analysis that compared the respiratory health of cement workers before and after the establishment of dust control measures, Tungu et al. (2014) [44] observed lower dust exposures among cement workers, lower prevalence of respiratory symptoms, chronic obstructive pulmonary disease, and improved lung function after introduction of dust control measures. Another study found a higher prevalence of respiratory symptoms among miners in a goldmine [45].

PAPERS

Three studies (one case-control and two cross-sectional) were conducted in Ethiopia between 2011 and 2023. All three focussed on the prevalence of respiratory diseases. Two examined dust exposure, while only one study investigated air pollution (PM_{2.5}, PM₁₀, CO₂, NO₂, SO₂) exposure. The results from the studies conducted in Ethiopia suggests higher level of dust exposure exceeding the local threshold limit of 10 mg/m³ for total dust and higher levels of air pollution (150 mg/m³ for particulate matter) [46,47]. Results from all studies conducted in Ethiopia found dust exposure and air pollution are linked with respiratory diseases [46–48]. Moreover, one cross-sectional study conducted in Zambia suggest a link between ambient air pollution (PM_{2.5}, PM₁₀) and respiratory symptoms [49].

Two studies were conducted in Cameroon between 2016 and 2017. One found a median precipitation level of 154mm and a temperature of 77°F (25°C) were linked with the prevalence of diabetes [50]. Another study conducted in Cameroon using a time-series study design observed no association among temperature, rainfall, humidity, and influenza [51]. However, two studies conducted in Congo found air pollution (PM_{2.5}, NO₂, SO₂) is linked with prevalence and incidence of respiratory symptoms [52,53]. Another study conducted in Senegal also found a link between NO₂ exposure and incidence of acute respiratory infections after adjusting for relevant cofounders [54]. Meanwhile, one cross-sectional study in Senegal also found heavy metal exposure is linked with respiratory diseases [55].

Two studies were conducted in Malawi. In one of them, Nightingale et al. (2019) [56] found 48-hour of CO exposure was linked with prevalence of respiratory symptoms. However, a RCT in rural Malawi found no significant association between daily average CO exposure among children who experienced pneumonia and those who did not [57]. This null association may be due to the overall low level of CO in their assessment. In addition, this study did not assess PM, a common pollutant linked with respiratory diseases.

Although poor environmental conditions and exposures are linked with chronic diseases [9,21], certain health benefits may be derived from the natural environment such as exposure to greenery [58,59]. For example, a multi-country study conducted in Tanzania, Uganda, and South Africa among 1178 participants using the Normalized Difference Vegetation Index (NDVI) found that exposure to neighborhood greenness was associated with lower body mass index (BMI), lower odds of obesity (odds ratio (OR) = 0.73; 95% confidence interval (CI) = 0.62, 0.85) and lower odds of diabetes (OR = 0.77; 95% CI = 0.62, 0.96) [60].

Two studies conducted in Mozambique found temperature declines (2.4°C to 3.0°C or 36.32°F to 37.4°F) were associated with incidence of stroke hospitalisations [61,62]. In another study conducted in 21 countries of SSA, Cai et al. (2021) [63] found prior month average air pollution (PM_{2.5}) exposure was not associated with cough and acute lower respiratory infections among children under five years of age. This null association may be due to several reasons. First, a global PM_{2.5} model may not have captured local emissions that are linked with biomass fuel

burning in rural areas. Second, exposures were assigned at survey clusters of participants locations (coordinates were displaced by 2–10 km of participants residence). Third, self-reporting of respiratory disease outcomes may have led to recall bias.

In this review, 30 studies found an association between environmental conditions and various types of chronic diseases, suggesting a link between environmental conditions and NCDs, particularly respiratory diseases in SSA. Air pollution, particularly PM_{2.5}, was the most assessed environmental exposure among the included studies. Most of the studies examining the impact of air pollution on chronic diseases reported high levels of air pollution (PM_{2.5}) exceeding the World Health Organization's previous and current air quality

guidelines of 10 $\mu\text{g}/\text{m}^3$ and 5 $\mu\text{g}/\text{m}^3$ for $\text{PM}_{2.5}$ [64]. The highest range for $\text{PM}_{2.5}$ (120-973 $\mu\text{g}/\text{m}^3$) was reported in one study conducted in Ethiopia [46]. $\text{PM}_{2.5}$ is a common pollutant and is linked with respiratory symptoms, cancer, and cardiovascular diseases [9,63,65,66]. Air pollution is reported as a major cause of NCD mortality in SSA and was responsible for 1.1 million deaths across the region in 2019 [21].

Dust and meteorological variables were the other leading environmental conditions reported in our scoping review. The included studies suggest temperature declines are associated with stroke and occupational dust exposure is linked with respiratory symptoms in SSA. This suggests a need for increased environmental monitoring and health tracking programs in SSA. Although certain

SSA countries such as South Africa

and Tanzania have started implementing environmental control measures, there still exist higher levels of environmental pollution in many of SSA countries. Future investments might focus on cleaner energies, particularly in rural parts of SSA. There is also need for improved diseases surveillance and more comprehensive health-related databases in SSA.

PAPERS

Limitations

This review has some limitations. First, many of the included studies used self-reported diseases outcome, which may have increased the chances of recall bias, i.e. participants may not have remembered important details regarding symptoms of past diseases. Second, exposure misclassification may have occurred; many of the studies assessing certain environmental exposures such as air pollution assume the variables measured at a few sites represented exposure to participants and for the larger geographic location. Third, we restricted this review to only studies published in English.

CONCLUSIONS

There is a modest, but growing body of research on environmental conditions and chronic diseases in SSA. This scoping review presented the existing evidence of the impact of environmental exposures on NCDs in SSA. Air pollution and respiratory diseases were the major environmental conditions and NCDs reported in studies assessed in this review. Our findings suggest an association between environmental conditions and NCDs, particularly respiratory diseases; we hope this study will provide guidance to researchers, policy makers, and international agencies to build upon existing research that will improve environmental health and reduce the burden of NCDs in SSA.

Acknowledgements: We would like to thank Rita Sieracki, reference librarian at the Medical College of Wisconsin Libraries, for her support with the electronic database searches.

Funding: None.

Authorship contributions: KB and CA conceived the study design. CA supported the systematic search, and conducted the literature reviews, and data extraction. JCB was the second reviewer and supported the data extraction, editing, and literature reviews. CA drafted and edited the manuscripts. KB reviewed/edited the manuscript.

Disclosure of interest: The authors completed the ICMJE Unified Competing Interest Form (available upon request from the corresponding author) and disclose no relevant interests.

Additional material

2 Gouda HN, Charlson F, Sorsdahl K, Ahmadzada S, Ferrari AJ, Erskine H, et al. Burden of non-communicable diseases in sub-Saharan Africa, 1990–2017: results from the Global Burden of Disease Study 2017. *Lancet Glob Health*. 2019;7:e1375–87.

3 Holmes MD, Dalal S, Volmink J, Adebamowo CA, Njelekela M, Fawzi WW, et al. Non-communicable diseases in sub-Saharan Africa: the case for cohort studies. *PLoS Med*. 2010;7:e1000244.

4 Mudie K, Jin MM, Tan, Kendall L, Addo J, dos-Santos-Silva I, et al. Non-communicable diseases in sub-Saharan Africa: a scoping review of large cohort studies. *J Glob Health*. 2019;9:020409. 5 Dalal S, Beunza JJ, Volmink J, Adebamowo C, Bajunirwe F, Njelekela M, et al. Non-communicable diseases in sub-Saharan Africa: what we know now. *Int J Epidemiol*. 2011;40:885–901.

6 Opoku SK, Filho WL, Hubert F, Adejumo O. Climate Change and Health Preparedness in Africa: Analysing Trends in Six African Countries. *Int J Environ Res Public Health*. 2021;18:4672. 7 Suhr F, Steinert JI. Epidemiology of floods in sub-Saharan Africa: a systematic review of health outcomes. *BMC Public Health*. 2022;22:268.

REFERENCES

9 Rodopoulou S, Samoli E, Chalbot MG, Kavouras IG. Air pollution and cardiovascular and respiratory emergency visits in Central Arkansas: A time-series analysis. *Sci Total Environ*. 2015;536:872-9.

10 Coleman NC, Burnett RT, Higbee JD, Lefler JS, Merrill RM, Ezzati M, et al. Cancer mortality risk, fine particulate air pollution, and smoking in a large, representative cohort of US adults. *Cancer Causes Control*. 2020;31:767-76.

11 Carpenter DO, Ma J, Lessner L. Asthma and infectious respiratory disease in relation to residence near hazardous waste sites. *Ann N Y Acad Sci*. 2008;1140:201-8.

12 Kudyakov R, Baibergenova A, Zdeb M, Carpenter DO. Respiratory disease in relation to patient residence near to hazardous waste sites. *Environ Toxicol Pharmacol*. 2004;18:249-57. 13 Kousky C, Kunreuther H, LaCour-Little M, Wachter S. Flood Risk and the U.S. Housing Market. *J Hous Res*. 2020;29:S3-24.

14 Tate E, Rahman MA, Emrich CT, Sampson CC. Flood exposure and social vulnerability in the United States. *Nat Hazards*. 2021;106:435-57.

15 Li Z, Shi J, Li N, Wang M, Jin Y, Zheng Z. Temporal trends in the burden of non-communicable diseases in countries with the highest malaria burden, 1990-2019: Evaluating the double burden of non-communicable and communicable diseases in epidemiological transition. *Global Health*. 2022;18:90. 16 Brainin P, Mohr GH, Modin D, Claggett B, Silvestre OM, Shah A, et al. Heart failure associated with imported malaria: a nationwide Danish cohort study. *ESC Heart Fail*. 2021;8:3521-9.

17 Mruma HA, McQuillan R, Norrie J. The association of malaria infection and gestational hypertension in Africa: Systematic review and meta-analysis. *J Glob Health*. 2020;10:020417.

18 Coogan PF, White LF, Jerrett M, Brook RD, Su JG, Seto E, et al. Air Pollution and Incidence of Hypertension and Diabetes in African American Women Living in Los Angeles. *Circulation*. 2012;125:767-72.

19 Ekolu J, Dieppois B, Sidibe M, Eden JM, Trambly Y, Villarini G, et al. Long-term variability in hydrological droughts and floods in sub-Saharan Africa: New perspectives from a 65-year daily streamflow dataset. *J Hydrol (Amst)*. 2022;613:128359.

20 Laraia BA. Food Insecurity and Chronic Disease. *Adv Nutr*. 2013;4:203-12.

21 Fisher S, Bellinger DC, Cropper ML, Kumar P, Binagwaho A, Koudenoukpo JB, et al. Air pollution and development in Africa: impacts on health, the economy, and human capital. *Lancet Planet Health*. 2021;5:e681-8.

22 Peters MDJ, Godfrey C, McInerney P, Khalil H, Larsen P, Marnie C, et al. Best practice guidance and reporting items for the development of scoping review protocols. *JBI Evid Synth*. 2022;20:953. 23 Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. *Syst Rev*. 2016;5:210.

24 Nang C, Piano B, Lewis A, Lycett K, Woodhouse M. Using The PICOS Model To Design And Conduct A Systematic Search: A Speech Pathology Case Study. Available: <https://ro.ecu.edu.au/cgi/viewcontent.cgi?article=1010&context=ecupres&httpsredir=1&referer=>. Accessed: 20 February 2024.

25 Norman R, Cairncross E, Witi J, Bradshaw D. South African Comparative Risk Assessment Collaborating G. Estimating the burden of disease attributable to urban outdoor air pollution in South Africa in 2000. *S Afr Med J*. 2007;97:782-90.

26 Kistnasamy EJ, Robins TG, Naidoo R, Batterman S, Mentz GB, Jack C, et al. The relationship between asthma and ambient air pollutants among primary school students in Durban, South Africa. *Int J Environ Health*. 2008;2:365-85.

27 Wichmann J, Vuyi K. Ambient air pollution exposure and respiratory, cardiovascular and cerebrovascular mortality in Cape Town, South Africa: 2001-2006. *Int J Environ Res Public Health*. 2012;9:3978-4016.

28 Naidoo RN, Robins TG, Batterman S, Mentz G, Jack C. Ambient pollution and respiratory outcomes among schoolchildren in Durban, South Africa. *SAJCH*. 2013;7:127-34.

29 Olaniyan T, Jeebhay M, Roosli M, Naidoo RN, Kunzli N, de Hoogh K, et al. The association between ambient NO₂ and PM_{2.5} with the respiratory health of school children residing in informal settlements: A prospective cohort study. *Environ Res*. 2020;186:109606.

30 Thabethe ND, Vuyi K, Wichmann J. Association between ambient air pollution and cause-specific mortality in Cape Town, Durban, and Johannesburg, South Africa: any susceptible groups? *Environ Sci Pollut Res Int*. 2021;28:42868-76.

PAPERS

31 Adebayo-Ojo TC, Wichmann J, Arowosegbe OO, Probst-Hensch N, Schindler C, Kunzli N. Short-Term Joint Effects of

PM10, NO2 and SO2 on Cardio-Respiratory Disease Hospital Admissions in Cape Town, South Africa. *Int J Environ Res Public Health*. 2022;19:495.

32 Shirinde J, Wichmann J. Temperature modifies the association between air pollution and respiratory disease mortality in Cape Town, South Africa. *Int J Environ Health Res*. 2023;33:1122–1131.

33 Wyndham CH. Are mortality rates for respiratory diseases in the RSA affected by climate? *S Afr Med J*. 1986;69:223–6.

34 Hnizdo E, Sluis-Cremer GK. Silica exposure, silicosis, and lung cancer: a mortality study of South African gold miners. *Br J Ind Med*. 1991;48:53–60.

PAPERS

35 Churchyard GJ, Ehrlich R, teWaterNaude JM, Pemba L, Dekker K, Vermeijs M, et al. Silicosis prevalence and exposure-response relations in South African goldminers. *Occup Environ Med*. 2004;61:811–6.

36 Millar DA, Kapwata T, Kunene Z, Mogotsi M, Wernecke B, Garland RM, et al. Respiratory health among adolescents liv

ing in the Highveld Air Pollution Priority Area in South Africa. *BMC Public Health*. 2022;22:2136.

37 Eghomwanre AF, Oguntoké O, Taiwo AM. Levels of indoor particulate matter and association with asthma in children in Benin City, Nigeria. *Environ Monit Assess*. 2022;194:467. 38 Oloyede T, Akintunde AA, Oloyede TW, Adeniran JA, Desalu OO, Tanimowo MO, et al. Air Quality Index and Respiratory

Symptoms among Garri Processing Workers in Ogbomoso, Nigeria. *West Afr J Med*. 2020;37:152–8. 39 Aliyu YA, Botai JO. An Exposure Appraisal of Outdoor Air Pollution on the Respiratory Well-being of a Developing City Population. *J Epidemiol Glob Health*. 2018;8:91–100.

40 Eghomwanre AF, Oguntoké O. Concentrations of indoor gaseous air pollutants and risk factors associated with childhood asthma in Benin City, Nigeria. *Environ Monit Assess*. 2022;194:391. 41 Farai IP, Obed RI, Jibiri NN. Soil radioactivity and incidence of cancer in Nigeria. *J Environ Radioact*. 2006;90:29–36.

42 Kilabuko JH, Matsuki H, Nakai S. Air quality and acute respiratory illness in biomass fuel using homes in Bagamoyo, Tanzania. *Int J Environ Res Public Health*. 2007;4:39–44.

43 Sakwari G, Bratveit M, Mamuya SH, Moen BE. Dust exposure and chronic respiratory symptoms among coffee curing workers in Kilimanjaro: a cross sectional study. *BMC Pulm Med*. 2011;11:54. 44 Tungu AM, Bratveit M, Mamuya SH, Moen BE. The impact of reduced dust exposure on respiratory health among cement workers: an ecological study. *J Occup Environ Med*. 2014;56:101–10. 45 Rusibamayila M, Meshi E, Mamuya S. Respiratory Impairment and Personal Respirable Dust Exposure among the Underground and Open Cast Gold Miners in Tanzania. *Ann Glob Health*. 2018;84:419–28.

46 Beketie KT, Angessa AT, Zeleke TT, Ayal DY. Impact of cement factory emission on air quality and human health around Mugher and the surrounding villages, Central Ethiopia. *Air Qual Atmos Health*. 2022;15:347–61.

47 Negash BM, Abaya SW, Abegaz T, Takele AK, Mekonnen WT, Negatu HB, et al. Assessment of paper dust exposure and chronic respiratory symptoms among paper factory workers in Ethiopia; a comparative cross-sectional study. *BMC Pulm Med*. 2023;23:48.

48 Zeleke ZK, Moen BE, Bratveit M. Lung function reduction and chronic respiratory symptoms among workers in the cement industry: a follow up study. *BMC Pulm Med*. 2011;11:50.

49 Nkhama E, Ndhlovu M, Dvonch JT, Lynam M, Mentz G, Siziya S, et al. Effects of Airborne Particulate Matter on Respiratory Health in a Community near a Cement Factory in Chilanga, Zambia: Results from a Panel Study. *Int J Environ Res Public Health*. 2017;14:1351.

50 Lontchi-Yimagou E, Tsalefac M, Tapinmene LM, Noubiap JJ, Balti EV, Nguewa JL, et al. Seasonality in diabetes in Yaounde, Cameroon: a relation with precipitation and temperature. *BMC Public Health*. 2016;16:470.

51 Monamele GC, Vernet MA, Nsaibirni RFJ, Bigna JJR, Kenmoe S, Njankouo MR, et al. Associations between meteorological parameters and influenza activity in a subtropical country: Case of five sentinel sites in Yaounde-Cameroon. *PLoS One*. 2017;12:e0186914.

52 Mbelambela EP, Hirota R, Eitoku M, Muchanga SMJ, Kiyosawa H, Yasumitsu-Lovell K, et al. Occupation exposed to road-traffic emissions and respiratory health among Congolese transit workers, particularly bus conductors, in Kinshasa: a cross-sectional study. *Environ Health Prev Med*. 2017;22:11. 53 Michellier C, Katoto PMC, Dramaix M, Nemery B, Kervyn F. Respiratory health and eruptions of the Nyiragongo and Nyamulagira volcanoes in the Democratic Republic of Congo: a time-series analysis. *Environmental Health:*

A Global Access Science Source. 2020;19:62.

54 Kawano A, Kim Y, Meas M, Sokal-Gutierrez K. Association between satellite-detected tropospheric nitrogen dioxide and acute respiratory infections in children under age five in Senegal: spatio-temporal analysis. BMC Public Health. 2022;22:178.

55 Faomowe Foko R, Mbodji M, Kêdote M, Diaw M, Touré M, Bah F, et al. Chemical risk and respiratory health of waste electrical and electronic equipment (WEEE) handlers in Dakar (Senegal). Arch Mal Prof Environ. 2021;82:438–48

56 Nightingale R, Lesosky M, Flitz G, Rylance SJ, Meghji J, Burney P, et al. Noncommunicable Respiratory Disease and Air Pollution Exposure in Malawi (CAPS). A Cross-Sectional Study. *Am J Respir Crit Care Med*. 2019;199:613–21.

REFERENCES

57 Mortimer K, Lesosky M, Semple S, Malava J, Katundu C, Crampin A, et al. Pneumonia and Exposure to Household Air Pollution in Children Under the Age of 5 Years in Rural Malawi: Findings From the Cooking and Pneumonia Study. *Chest*. 2020;158:501–11.

58 Barton J, Rogerson M. The importance of greenspace for mental health. *BJPsych Int*. 2017;14:79–81.

59 Bikomeye JC, Namin S, Anyanwu C, Rublee CS, Ferschinger J, Leinbach K, et al. Resilience and Equity in a Time of Crises: Investing in Public Urban Greenspace Is Now More Essential Than Ever in the US and Beyond. *Int J Environ Res Public Health*. 2021;18:8420.

60 Iyer HS, James P, Valeri L, Bajunirwe F, Nankya-Mutyoba J, Njelekela M, et al. Neighborhood greenness and burden of non-communicable diseases in Sub-Saharan Africa: A multi-country cross-sectional study. *Environ Res*. 2021;196:110397.

PAPERS

61 Gomes J, Damasceno A, Carrilho C, Lobo V, Lopes H, Madede T, et al. Triggering of stroke by ambient temperature variation: a case-crossover study in Maputo, Mozambique. *Clin Neurol Neurosurg*. 2015;129:72–7.

62 Gomes J, Damasceno A, Carrilho C, Lobo V, Lopes H, Madede T, et al. The effect of season and temperature variation on hospital admissions for incident stroke events in Maputo, Mozambique. *J Stroke Cerebrovasc Dis*. 2014;23:271–7.

63 Cai YS, Gibson H, Ramakrishnan R, Mamouei M, Rahimi K. Ambient air pollution and respiratory health in sub-Saharan African children: A cross-sectional analysis. *Int J Environ Res Public Health*. 2021;18:9729.

64 World Health Organization. What are the WHO Air quality guidelines? Available: . Accessed: 26 September 2023.

65 Servadio JL, Lawal AS, Davis T, Bates J, Russell AG, Ramaswami A, et al. Demographic Inequities in Health Outcomes and Air Pollution Exposure in the Atlanta Area and its Relationship to Urban Infrastructure. *J Urban Health*. 2019;96:219–34.

66 Turner MC, Andersen ZJ, Baccarelli A, Diver WR, Gapstur SM, Pope CA III, et al. Outdoor air pollution and cancer: An overview of the current evidence and public health recommendations. *CA Cancer J Clin*. 2020;70:460–79.

